

Renormalization theory of Herman curves

Willie Rush Lim

Brown University

Aug 4, 2025

Diophantine assumption

Fix an irrational $\theta \in (0, 1)$ and write

$$\theta = [a_1, a_2, a_3, \dots] := \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}.$$

θ is called

- **bounded** if $\sup a_n < \infty$.
- **periodic** with period p if $a_{n+p} = a_n$ for all n .

E.g. golden mean $= [1, 1, 1, \dots] = \frac{\sqrt{5}-1}{2}$

Diophantine assumption

Fix an irrational $\theta \in (0, 1)$ and write

$$\theta = [a_1, a_2, a_3, \dots] := \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}}.$$

θ is called

- **bounded** if $\sup a_n < \infty$.
- **periodic** with period p if $a_{n+p} = a_n$ for all n .

E.g. golden mean $= [1, 1, 1, \dots] = \frac{\sqrt{5}-1}{2}$

The n^{th} rational approximation of θ is

$$\frac{p_n}{q_n} = [a_1, \dots, a_n].$$

Rotation curves

An invariant Jordan curve $\mathbf{H} \subset \hat{\mathbb{C}}$ of a holomorphic map f is

- a **rotation curve** if $f|_{\mathbf{H}}$ is conjugate to an irrational rotation $R_\theta : \mathbb{T} \rightarrow \mathbb{T}$;
- a **Herman curve** if additionally it isn't contained in the closure of a rotation domain.

Rotation curves

An invariant Jordan curve $\mathbf{H} \subset \hat{\mathbb{C}}$ of a holomorphic map f is

- a **rotation curve** if $f|_{\mathbf{H}}$ is conjugate to an irrational rotation $R_\theta : \mathbb{T} \rightarrow \mathbb{T}$;
- a **Herman curve** if additionally it isn't contained in the closure of a rotation domain.

Trichotomy: When a rotation curve \mathbf{H} has bounded rotation number θ , there are 3 cases:

- \mathbf{H} = an analytic curve inside a rotation domain,
- \mathbf{H} = the boundary of a rotation domain containing a critical point of f ,
- \mathbf{H} = a Herman curve containing inner and outer critical points of f .



Rotation curves

An invariant Jordan curve $\mathbf{H} \subset \hat{\mathbb{C}}$ of a holomorphic map f is

- a **rotation curve** if $f|_{\mathbf{H}}$ is conjugate to an irrational rotation $R_{\theta} : \mathbb{T} \rightarrow \mathbb{T}$;
- a **Herman curve** if additionally it isn't contained in the closure of a rotation domain.

Trichotomy: When a rotation curve \mathbf{H} has bounded rotation number θ , there are 3 cases:

- \mathbf{H} = an analytic curve inside a rotation domain,
- \mathbf{H} = the boundary of a rotation domain containing a critical point of f ,
- \mathbf{H} = a Herman curve containing inner and outer critical points of f .



Siegel disks and Herman rings have been pretty well studied by many many people. On the other hand, not much is known about Herman curves.

Key questions

- 1 Regularity and smoothness of Herman curves?
- 2 Rigidity properties?
- 3 Regularity of conjugacy classes?
- 4 Structural instability?

From now on, we will consider a Herman curve $f : \mathbb{H} \rightarrow \mathbb{H}$ with a single critical point c .


Inner & outer criticalities

Denote

$d_0 =$ inner criticality of c ,

$d_\infty =$ outer criticality of c .

The total local degree of the critical point c is $d_0 + d_\infty - 1$.



Example of
 $(d_0, d_\infty) = (2, 3)$


Inner & outer criticalities

Denote

$d_0 =$ inner criticality of c ,

$d_\infty =$ outer criticality of c .

The total local degree of the critical point c is $d_0 + d_\infty - 1$.



Critical circle maps (when $\mathbf{H} = S^1$) automatically have $d_0 = d_\infty$.


E.g. The Arnold family, $(d_0, d_\infty) = (2, 2)$:

$$A_t(x) = x + t - \frac{1}{2\pi} \sin(2\pi x), \quad x \in \mathbb{R}/\mathbb{Z}.$$

Blaschke product example, $d_0 = d_\infty = 2$

For any irrational θ , there is a unique $c_\theta \in \mathbb{T}$ such that the unit circle is a Herman curve of rotation number θ for the map

$$f_\theta(z) = c_\theta z^2 \frac{z-3}{1-3z}.$$



Can we generalize this?

Arbitrary criticalities


Fix a bounded irrational θ , a pair (d_0, d_∞) , and $d := d_0 + d_\infty - 1$.

Realization+Uniqueness Theorem [wrl '23]

There exists a unique degree d rational map

$$F : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$$

that has the critical portrait below and a Herman quasicycle \mathbf{H} with rotation number θ .



Arbitrary criticalities


Fix a bounded irrational θ , a pair (d_0, d_∞) , and $d := d_0 + d_\infty - 1$.

Realization+Uniqueness Theorem [wrl '23]

There exists a unique degree d rational map


$$F : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$$

that has the critical portrait below and a Herman quasicycle \mathbf{H} with rotation number θ .



The realization follows from a priori bounds and degeneration of Herman rings. The uniqueness follows from the absence of line fields in the Julia set.

Example for $(d_0, d_\infty) = (3, 2)$



$\theta = \text{golden mean}$


$$F_{c_*}(z) = c_* z^3 \frac{4 - z}{1 - 4z + 6z^2}$$

$$c_* \approx -1.14421 - 0.96445i$$

The map F_{c_*} naturally lives in the 1-parameter family

$$\left\{ F_c = cz^3 \frac{4 - z}{1 - 4z + 6z^2} \right\}_{c \in \mathbb{C}^*}.$$

The parameter space picture



Conjecture: The bifurcation locus of $\{F_c\}_{c \in \mathbb{C}^*}$ is asymptotically self-similar at c_* .

Going beyond rational maps

From now on, fix integers $d_0, d_\infty \geq 2$. Denote the set

$$\mathrm{HC}_\theta = \left\{ (f, \mathbf{H}) : \begin{array}{l} f \text{ is a holomorphic map and } \mathbf{H} \text{ is a} \\ \text{unicritical Herman quasicircle of } f \text{ with} \\ \text{rotation number } \theta \text{ and criticalities } (d_0, d_\infty) \end{array} \right\}.$$

Going beyond rational maps

From now on, fix integers $d_0, d_\infty \geq 2$. Denote the set

$$\mathrm{HC}_\theta = \left\{ (f, \mathbf{H}) : \begin{array}{l} f \text{ is a holomorphic map and } \mathbf{H} \text{ is a} \\ \text{unicritical Herman quasicycle of } f \text{ with} \\ \text{rotation number } \theta \text{ and criticalities } (d_0, d_\infty) \end{array} \right\}.$$

Rigidity Theorem [wrl '23]

For any bounded type θ and any (f, \mathbf{H}) and $(\tilde{f}, \tilde{\mathbf{H}})$ in HC_θ ,

- ① there is a qc conjugacy ϕ between f and \tilde{f} on a nbh of \mathbf{H} ,
- ② ϕ is $C^{1+\alpha}$ -conformal on \mathbf{H} .

In the special case $\mathbf{H} = \tilde{\mathbf{H}} = S^1$, this was proven by de Faria-de Melo '00.

Going beyond rational maps

From now on, fix integers $d_0, d_\infty \geq 2$. Denote the set

$$\mathrm{HC}_\theta = \left\{ (f, \mathbf{H}) : \begin{array}{l} f \text{ is a holomorphic map and } \mathbf{H} \text{ is a} \\ \text{unicritical Herman quasicycle of } f \text{ with} \\ \text{rotation number } \theta \text{ and criticalities } (d_0, d_\infty) \end{array} \right\}.$$

Rigidity Theorem [wrl '23]

For any bounded type θ and any (f, \mathbf{H}) and $(\tilde{f}, \tilde{\mathbf{H}})$ in HC_θ ,

- 1 there is a qc conjugacy ϕ between f and \tilde{f} on a nbh of \mathbf{H} ,
- 2 ϕ is $C^{1+\alpha}$ -conformal on \mathbf{H} .

In the special case $\mathbf{H} = \tilde{\mathbf{H}} = S^1$, this was proven by de Faria-de Melo '00.

Rigidity has many consequences, e.g.

- 1 $\dim(\mathbf{H}) = \text{universal constant}$;
- 2 $\dim(\mathbf{H}) = 1 \iff \mathbf{H} \text{ is } C^1\text{-smooth} \iff d_0 = d_\infty$;
- 3 if θ is pre-periodic, \mathbf{H} is self-similar at the crit. pt. with universal scaling constant.

Renormalization


Denote $\{c_i := f^i(c)\}_{i \geq 0}$ the critical orbit of f .

The n^{th} **pre-renormalization** $p\mathcal{R}^n f$ is the pair

$$\left(f^{q_n}|_{[c_{q_{n-1}}, c_0]}, f^{q_{n-1}}|_{[c_0, c_{q_n}]} \right)$$

which is the first return map of f back to the interval $[c_{q_{n-1}}, c_{q_n}] \subset \mathbf{H}$.

The n^{th} **renormalization** $\mathcal{R}^n f$ is the normalized pair obtained by affine rescaling $c_{q_{n-1}} \mapsto -1$ and $c_0 \mapsto 0$.



Renormalization


Denote $\{c_i := f^i(c)\}_{i \geq 0} =$ the critical orbit of f .

The **n^{th} pre-renormalization** $p\mathcal{R}^n f$ is the pair

$$\left(f^{q_n}|_{[c_{q_{n-1}}, c_0]}, f^{q_{n-1}}|_{[c_0, c_{q_n}]} \right)$$

which is the first return map of f back to the interval $[c_{q_{n-1}}, c_{q_n}] \subset \mathbf{H}$.

The **n^{th} renormalization** $\mathcal{R}^n f$ is the normalized pair obtained by affine rescaling $c_{q_{n-1}} \mapsto -1$ and $c_0 \mapsto 0$.




\mathcal{R} acts on rotation number as the Gauss map:

$$\text{rot}(f) = \theta = [a_1, a_2, \dots] \implies \text{rot}(\mathcal{R}^n f) = G^n \theta = [a_{n+1}, a_{n+2}, \dots].$$



Outline of proof of Rigidity

1. Petersen '04: $f|_{\mathbf{H}} \sim_{\text{qs}} \tilde{f}|_{\tilde{\mathbf{H}}}$
2. Prove uniform butterfly structure (complex bounds) for $p\mathcal{R}^n f$. $n \gg 1$.
3. Construct qc conjugacy between the butterflies of $p\mathcal{R}^n f$ and $p\mathcal{R}^n \tilde{f}$.
4. Spread this around to get a conjugacy ϕ on a nbh of \mathbf{H} .



Outline of proof of Rigidity

1. Petersen '04: $f|_{\mathbf{H}} \sim_{\text{qs}} \tilde{f}|_{\tilde{\mathbf{H}}}$
2. Prove uniform butterfly structure (complex bounds) for $p\mathcal{R}^n f$. $n \gg 1$.
3. Construct qc conjugacy between the butterflies of $p\mathcal{R}^n f$ and $p\mathcal{R}^n \tilde{f}$.
4. Spread this around to get a conjugacy ϕ on a nbh of \mathbf{H} .
5. Show that $\bar{\partial}\phi = 0$ a.e. on $J_f = \overline{\cup_{k \geq 0} f^{-k}(\mathbf{H})}$.
6. Prove that points on \mathbf{H} are “uniformly deep” in J_f :



As we zoom in near the critical pt, J_f converges to \mathbb{C} exp. fast.

Renormalization fixed point

Fix a periodic irrational θ_* with some even period p .

Corollary

There is a unique normalized commuting pair ζ_ with rot. no. θ_* satisfying $\mathcal{R}^p \zeta_* = \zeta_*$.
For any $(f, \mathbf{H}) \in \text{HC}_{\theta_*}$,*

$$\mathcal{R}^{np} f \longrightarrow \zeta_* \quad \text{exp. fast as } n \rightarrow \infty.$$


Renormalization fixed point

Fix a periodic irrational θ_* with some even period p .

Corollary

There is a unique normalized commuting pair ζ_* with rot. no. θ_* satisfying $\mathcal{R}^p \zeta_* = \zeta_*$.
For any $(f, \mathbf{H}) \in \text{HC}_{\theta_*}$,

$$\mathcal{R}^{np} f \longrightarrow \zeta_* \quad \text{exp. fast as } n \rightarrow \infty.$$



Renormalization fixed point


Fix a periodic irrational θ_* with some even period p .

Corollary

There is a unique normalized commuting pair ζ_* with rot. no. θ_* satisfying $\mathcal{R}^p \zeta_* = \zeta_*$.
For any $(f, \mathbf{H}) \in \text{HC}_{\theta_*}$,

$$\mathcal{R}^{np} f \longrightarrow \zeta_* \quad \text{exp. fast as } n \rightarrow \infty.$$

One can also glue the two ends of the commuting pair ζ_* to obtain a Herman quasicircle $f_* : \mathbf{H}_* \rightarrow \mathbf{H}_*$ fixed by a renormalization operator \mathcal{R}_* :



Fix a skinny annular nbh A of \mathbf{H}_* and a small $\varepsilon > 0$. Define the Banach ball:

$$\mathcal{B}_\varepsilon(f_*) := \left\{ g \in \text{Hol}(A, \mathbb{C}) \mid g \text{ has a unique critical point and } \sup_{z \in A} |g(z) - f(z)| < \varepsilon \right\}.$$

Hyperbolicity of renormalization

Fix a skinny annular nbh A of \mathbf{H}_* and a small $\varepsilon > 0$. Define the Banach ball:

$$\mathcal{B}_\varepsilon(f_*) := \left\{ g \in \text{Hol}(A, \mathbb{C}) \mid g \text{ has a unique critical point and } \sup_{z \in A} |g(z) - f(z)| < \varepsilon \right\}.$$

Hyperbolicity Theorem [wrl '24]


\mathcal{R}_* can be naturally extended to a compact analytic operator on $\mathcal{B}_\varepsilon(f_*)$ such that:

- ① f_* is the unique fixed point of \mathcal{R}_* .
- ② f_* is hyperbolic with a single unstable direction.
- ③ $\mathcal{W}_{\text{loc}}^s(f_*) = \{g \in \mathcal{B}_\varepsilon(f_*) \mid g \text{ has a Herman quasicircle with rot. no. } \theta_*\}.$

In the circle case ($d_0 = d_\infty$), the real version of this was proven by Yampolsky '03.

Key ingredient: Corona structure


A **corona** is a holomorphic map $f : U \rightarrow V$ between nested annuli with radial arcs $\gamma_0 \subset U$ and $\gamma_1 \subset V$ such that $f : U \setminus \gamma_0 \rightarrow V \setminus \gamma_1$ is a covering map branched at a unique crit. pt.



A corona $f : (U, \gamma_0) \rightarrow (V, \gamma_1)$ with criticalities $(d_0, d_\infty) = (2, 3)$


Corona renormalization operator

Every unicritical Herman curve can be renormalized to a corona:



Corona renormalization operator

Every unicritical Herman curve can be renormalized to a corona:



\mathcal{R}_{cor} naturally extends to an analytic operator on a Banach ball $\mathcal{B}_\varepsilon(f)$.

Since $f_* : \mathbf{H}_* \rightarrow \mathbf{H}_*$ can be renormalized to itself, f_* admits a corona structure. We extend $\mathcal{R}_* : f_* \mapsto f_*$ to an analytic renormalization operator on $\mathcal{B}_\varepsilon(f_*)$.

Most difficult part of the proof

With this corona framework, together with various soft methods¹, most of the hyperbolicity theorem can be proven.

Remaining obstacle: $\dim(\mathcal{W}_{\text{loc}}^u) \leq 1$?

¹holomorphic motions, renormalization tiling, Small Orbits Theorem, exponential convergence, etc

Most difficult part of the proof

With this corona framework, together with various soft methods¹, most of the hyperbolicity theorem can be proven.

Remaining obstacle: $\dim(\mathcal{W}_{\text{loc}}^u) \leq 1$?

Solution: $\mathcal{W}_{\text{loc}}^u$ = a parameter space of transcendental maps of unknown dimension.

¹holomorphic motions, renormalization tiling, Small Orbits Theorem, exponential convergence, etc

Transcendental dynamics

For the renormalization fixed point f_* ,

$$\begin{array}{ccc} \text{rescalings of} & & \text{bi-infinite tower of commuting } \sigma\text{-proper maps} \\ \left\{ f_*^{Q_P(m+n)} \right\}_{m \geq -n} & \xrightarrow{m \rightarrow \infty} & \left\{ \mathbf{F}_*^{Q_{pn}} : \text{Dom}(\mathbf{F}_*^{Q_{pn}}) \rightarrow \mathbb{C} \right\}_{n \in \mathbb{Z}} \end{array}$$

We can normalize this process so that 0 is the critical value.

$\left\{ \mathbf{F}_*^{Q_n} \right\}_{n \in \mathbb{Z}}$ generates a semigroup of transcendental maps $\mathbf{F}_* = (\mathbf{F}_*^P)_{P \in \mathbf{T}}$.

Transcendental dynamics

For the renormalization fixed point f_* ,

$$\begin{array}{ccc} \text{rescalings of} & & \text{bi-infinite tower of commuting } \sigma\text{-proper maps} \\ \left\{ f_*^{Q_P(m+n)} \right\}_{m \geq -n} & \xrightarrow{m \rightarrow \infty} & \left\{ \mathbf{F}_*^{Q_{pn}} : \text{Dom}(\mathbf{F}_*^{Q_{pn}}) \rightarrow \mathbb{C} \right\}_{n \in \mathbb{Z}} \end{array}$$

We can normalize this process so that 0 is the critical value.

$$\left\{ \mathbf{F}_*^{Q_n} \right\}_{n \in \mathbb{Z}} \text{ generates a semigroup of transcendental maps } \mathbf{F}_* = (\mathbf{F}_*^P)_{P \in \mathbf{T}}.$$

For $f \in \mathcal{W}_{\text{loc}}^u$,

$$\begin{array}{ccc} \left\{ \mathcal{R}^{-n} f \right\}_{n \leq 0} & \xrightarrow[\text{rescaling}]{\text{appropriate}} & \text{backward tower of commuting } \sigma\text{-proper maps} \\ & & \left\{ \mathbf{F}^{Q_{pn}} : \text{Dom}(\mathbf{F}^{Q_{pn}}) \rightarrow \mathbb{C} \right\}_{n \leq 0} \end{array}$$

This still forms a semigroup of transcendental maps $\mathbf{F} = (\mathbf{F}^P)_{P \in \mathbf{T}}$.

Dynamical sets for cascades

For $f \in \mathcal{W}^u$, define...

- Fatou set:

$$\mathfrak{F}(\mathbf{F}) = \text{points of normality of } (\mathbf{F}^P)_{P \in \mathbf{T}}$$

- Julia set:

$$\mathfrak{J}(\mathbf{F}) = \mathbb{C} \setminus \mathfrak{F}(\mathbf{F})$$

- postcritical set:

$$\mathfrak{P}(\mathbf{F}) = \text{closure of the critical orbit } (\mathbf{F}^P(0))_{P \in \mathbf{T}}$$

- finite-time escaping set:

$$\mathbf{I}_{<\infty}(\mathbf{F}) = \bigcup_{P \in \mathbf{T}} \mathbb{C} \setminus \text{Dom}(\mathbf{F}^P)$$


- infinite-time escaping set:

$$\mathbf{I}_{\infty}(\mathbf{F}) = \text{points } x \text{ where } \mathbf{F}^P(x) \rightarrow \infty \text{ as } P \rightarrow \infty$$

- full escaping set:

$$\mathbf{I}(\mathbf{F}) = \mathbf{I}_{<\infty}(\mathbf{F}) \cup \mathbf{I}_{\infty}(\mathbf{F}).$$

Approximate dynamical picture of \mathbf{F}_*



In blue:
Some rays in $\mathbf{I}_{<\infty}(\mathbf{F}_*)$
landing at critical points of \mathbf{F}_*

$\mathfrak{P}(\mathbf{F}_*)$

\mathcal{W}^u is one-dimensional

Rigidity Theorem for \mathcal{W}^u

For $f \in \mathcal{W}_{\text{loc}}^n$,

1. $\mathbf{I}(\mathbf{F})$ supports no invariant line field & moves conformally away from pre-critical pts.
2. If \mathbf{F} is hyperbolic, then $\mathfrak{J}(\mathbf{F})$ also supports no invariant line field.

At last,

$$\begin{array}{c} \exists \text{ hyperbolic component} \\ O \subset \mathcal{W}_{\text{loc}}^u \text{ near } f_* \end{array} \xrightarrow[\text{above}]{\text{theorem}} \dim(O) \leq \# \text{ free critical orbits} = 1.$$

Structure of the local conjugacy class

Corollary

Consider any pre-periodic irrational θ' and any $(f, \mathbf{H}) \in \mathrm{HC}_{\theta'}$. The local conjugacy class

$$\left\{ g \in \mathcal{B}_\varepsilon(f) \mid g \text{ has a Herman quasicircle } \mathbf{H}_g \text{ with rot. no. } \theta' \right\}$$

is an analytic submanifold of $\mathcal{B}_\varepsilon(f)$ of $\mathrm{codim} \leq 1$ on which \mathbf{H}_g moves holomorphically.


Structure of the local conjugacy class

Corollary

Consider any pre-periodic irrational θ' and any $(f, \mathbf{H}) \in \text{HC}_{\theta'}$. The local conjugacy class

$$\left\{ g \in \mathcal{B}_\varepsilon(f) \mid g \text{ has a Herman quasicycle } \mathbf{H}_g \text{ with rot. no. } \theta' \right\}$$

is an analytic submanifold of $\mathcal{B}_\varepsilon(f)$ of $\text{codim} \leq 1$ on which \mathbf{H}_g moves holomorphically.



Development of renormalization theory in complex dynamics

real renorm. horseshoe	∞-renorm unimodal maps	critical circle maps
beau bounds	Sullivan, Lyubich-Yampolsky, Levin-van Strien	de Faria, Yampolsky, de Faria-de Melo
exp contraction	McMullen, Avila-Lyubich	de Faria-de Melo
hyperbolicity	Lyubich Avila-Lyubich	Yampolsky

renorm. fixed point	∞-renorm PL maps	Siegel disks	Herman curves
complex a priori bounds	Kahn, Dudko-Lyubich	McMullen, Avila-Lyubich	Lim
$C^{1+\alpha}$-rigidity/ exp contraction	McMullen, Lyubich	McMullen, Avila-Lyubich	Lim
hyperbolicity	Lyubich	Gaidashev-Yampolsky, Dudko-Lyubich-Selinger	Lim

The parameter picture

features	∞ -renorm. PL maps	Siegel disks	Herman curves
nice family	$z^d + c$	$z^d + c$	$-c \frac{\sum_{j=d_0}^d \binom{d}{j} \cdot (-z)^j}{\sum_{j=0}^{d_0-1} \binom{d}{j} \cdot (-z)^j}$
\mathcal{R} -invariant lamination	hybrid classes	level sets of multiplier	?
structural instability	yes	yes	partially known
parameter self-similarity	complete	partially known	?

Thank you!